

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

TABLE OF CONTENTS


Patient Initial Evaluation.....	Page 2
APPENDIX A: Checklist/Supportive Care Considerations for Managing Patients Receiving IEC Therapy...	Pages 3-5
APPENDIX B: Infectious Disease Screening.....	Page 6
APPENDIX C: Infection Prophylaxis Considerations for IEC Therapy.....	Pages 7-10
APPENDIX D: ASTCT Grading of CRS.....	Page 11
APPENDIX E: ASTCT Grading of ICANS.....	Page 12
APPENDIX F: Immune Effector Cell-associated Encephalopathy (ICE) Score.....	Page 13
APPENDIX G: Cornell Assessment of Pediatric Delirium (CAPD).....	Page 14
APPENDIX H: Management of CRS.....	Pages 15-17
APPENDIX I: Management of ICANS.....	Pages 18-20
APPENDIX J: Recommendations for Use of Interleukin-6 (IL-6) Antagonists and Alternative Agents for Management of CRS and ICANS.....	Page 21
APPENDIX K: Management of Focal or Generalized or Non-Convulsive Seizures.....	Page 22
APPENDIX L: Management of Convulsive Status Epilepticus.....	Page 22
APPENDIX M: Management of Diffuse Cerebral Edema and/or Raised Intracranial Pressure.....	Page 23
APPENDIX N: Diagnostic Criteria and Management for IEC-Related Hemophagocytic Lymphohistiocytosis (HLH) or Macrophage Activation Syndrome (MAS)	Page 24
APPENDIX O: Allogeneic IEC-associated Acute Graft-Versus-Host Disease (GVHD).....	Page 25
APPENDIX P: Grade of IEC-associated Acute GVHD.....	Page 25
APPENDIX Q: Manage IEC-associated Acute GVHD.....	Page 26
Suggested Readings.....	Page 27
Development Credits.....	Page 28

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

INITIAL EVALUATION

MANAGEMENT



¹ If the subject has fever with or without hypotension or hypoxia within the first 4 weeks of engineered immune effector cell (IEC) therapy, the subject may have CRS if the symptoms or signs are not attributable to any other cause:

- Fever should be present at onset of CRS (oral temperature $\geq 38^{\circ}\text{C}$)
- Hypotension (requiring IV fluids or vasopressors to maintain normal blood pressure) defined as:
 - Age 1-10 years: SBP < [70 + (2 x age in years)] mmHg
 - Age > 10 years: SBP < 90 mmHg
- Hypoxia (requiring supplemental oxygen to correct a deficit in oxygenation)

² If the subject has any of the following within the first 8 weeks of engineered IEC-therapy, the subject may have ICANS if the symptoms or signs are not attributable to any other cause:

- IEC-Associated Encephalopathy (ICE) Score of < 10 (Appendix F) and/or Cornell Assessment of Pediatric Delirium (CAPD) Score > 7 (see Appendix G)
- Depressed level of consciousness
- Convulsive or non-convulsive seizures (can be focal or generalized)
- Motor weakness (can be focal motor weakness, hemiparesis, paraparesis)
- Focal / diffuse cerebral edema on imaging or signs of raised intracranial pressure including decerebrate or decorticate posturing, cranial nerve VI palsy, papilledema, or Cushing's triad

³ See Appendix D for Grading of CRS

⁴ See Appendix E for Grading of ICANS

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX A: Checklist / Supportive Care Considerations for Managing Patients Receiving IEC Therapy

Before and During IEC Infusion

- Imaging of the brain prior to IEC infusion (preferably MRI with and without contrast but CT without contrast is acceptable if MRI cannot be performed) to rule out any central nervous system disease and also to serve as a baseline for comparison in case the patient develops ICANS
 - For patients with known history of seizures, migraines and/or other CNS disorders including malignant disease, consider Neurology consult prior to IEC infusion
- Central venous access with port-a-cath or double/triple lumen catheter is recommended for IEC infusion as well as for IV fluids and other infusions in case of toxicities
- IEC infusion may be administered either in the ambulatory unit or in the inpatient unit
- If the median time to onset of CRS is expected to be < 48 hours, hospitalization should be considered for IEC infusion
- When hospitalized, admission to an IEC-designated unit with capability for cardiac monitoring by telemetry is recommended
- Tumor lysis precautions for patients with high tumor burden, as per standard guidelines
- Seizure prophylaxis with levetiracetam 10 mg/kg (maximum 500 mg) PO or IV every 12 hours for 30 days, starting on the day of infusion for IEC therapies associated with a high incidence of ICANS, in patients with history of seizures or brain metastases
- Consider filgrastim products if patient is neutropenic and concern for infection (if not already receiving)
- Ensure appropriate documentation in EHR regarding IEC therapy and “conditional” corticosteroid contraindication

Continued on next page

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX A: Checklist / Supportive Care Considerations for Managing Patients Receiving IEC Therapy - *continued*

For Inpatients:

Patient Monitoring After IEC infusion during Inpatient Admission

- Assess vital signs every 4 hours (inpatient encounter)
- Strict monitoring of oral and IV fluid input and output (including urine and stool)
- Daily measurement of body weight
- Daily review of patient history and physical examination
- Daily complete blood count with differential and complete metabolic profile
- Coagulation profile at least twice weekly or more frequently if clinically indicated
- Consider monitoring C-reactive protein (CRP) and ferritin levels daily during the phase when CRS is likely to occur and continue to monitor until CRS and/or ICANS resolves (if present). Monitor as needed thereafter.
- Consider cytokine panel if clinically indicated
- Assessment and grading of CRS (document in CARTOX flowsheet) should be completed at least every 12 hours and whenever there is a change in patient's status. If a change in status occurs, monitor for CRS every 6 hours or more frequently as indicated.
- Assessment and grading for ICANS (document in CARTOX flowsheet) should be completed at least every 12 hours including the CAPD (if age or developmental age is ≤ 18 years of age) and 10-point ICE score if ≥ 12 years of age. If < 12 years of age, the CAPD should be utilized. Any increase in CAPD score requires monitoring every 6 hours or more frequently as indicated.
- Maintenance IV fluids with normal saline to ensure adequate hydration
- Cardiac monitoring by telemetry is recommended for \geq Grade 1 CRS and continued until CRS resolves
- For post-IEC infusion headache that is unresponsive to analgesics, consider brain imaging and lumbar puncture
- Neurology consult recommended for patients who develop Grade 1 or higher ICANS
- Critical Care and/or MERIT team will follow patients on an as-needed basis
- Infectious Diseases team will follow patients on an as-needed basis
 - Consult should be performed early for patients with positive infectious disease screening or for persistent fevers

Continued on next page

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX A: Checklist / Supportive Care Considerations for Managing Patients Receiving IEC Therapy- *continued*

Notifications and contingency orders during Inpatient Admission

- Notify SCT physician on detection of any of the following:

The baseline blood pressure should be considered to determine monitoring parameters and verified with the physician prior to admission

- Hypotension defined as:
 - Age 1-10 years: SBP < [70 + (2 x age in years)] mmHg
 - Age > 10 years: SBP < 90 mmHg
- Heart rate based on age or arrhythmia:
 - Age 1-2 years: > 130 or < 75 beats per minute
 - Age 3- 6 years: > 120 or < 70 beats per minute
 - Age 7-12 years: > 110 or < 65 beats per minute
 - Age 13-17 years: > 105 or < 55 beats per minute
- Respiratory rate based on age:
 - Age 1-2 years: > 40 or < 24 breaths per minute
 - Age 3- 6 years: > 34 or < 22 breaths per minute
 - Age 7-12 years: > 30 or < 18 breaths per minute
 - Age 13-17 years: > 16 or < 12 breaths per minute
- Oxygen saturation < 94% on room air
- Decreased urine output (<0.5 mL/kg/hour)
- Upward trends in creatinine or liver function tests
- Tremors or jerky movements in extremities
- Change in mental status (alertness, orientation, speech, ability to write a sentence, or ICE score of < 10 and/or CAPD score > 7)
- For oral temperature $\geq 38^{\circ}\text{C}$, send blood cultures (central and peripheral) and urine for urinalysis and culture, obtain portable chest x-ray, and notify physician
- For patients with neutropenia and fever, start empiric broad-spectrum antibiotics
- Do not administer corticosteroids unless approved by physician
- If patient develops ICANS, withhold oral intake of food, fluids, and medicines, and notify physician
- PRN medications
 - Acetaminophen (1st choice) or ibuprofen (2nd choice, if not contraindicated) for fever $\geq 38^{\circ}\text{C}$
 - Cooling blanket for fever $\geq 38.3^{\circ}\text{C}$
 - Normal saline 10 - 20 mL/kg (maximum 1,000 mL) bolus prn for hypotension; may repeat once if patient remains hypotensive after 1st bolus
 - Transfuse packed red blood cells (PRBC) to maintain hemoglobin > 8 gm/dL
 - Transfuse platelets to maintain > 20 K/microliter; for patients with abnormal brain imaging transfuse platelets to maintain > 50 k/microliter
 - PRN tocilizumab to be activated only on physician order (“ok to give tocilizumab” order should be placed if dose approved by physician)

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX B: Infectious Disease Screening (within 30 days prior to apheresis is recommended)

<p>Required Infectious Disease Screening¹</p> <ul style="list-style-type: none"> • Hepatitis B surface antigen (HBsAg) • Anti-hepatitis B core antibody (HBcAb) • Anti-hepatitis C virus antibody (HCVAb) • Anti-human immunodeficiency virus (HIV) antibody (HIV type 1 / 2 type O Ab) • HIV-1 / HCV / HBV Nucleic Acid Test • HHV-6 IgG (Herpesvirus 6 Ab panel) • Cytomegalovirus (CMV) IgG and IgM 	<p>Optional Infectious Disease Screening (as clinically indicated)²</p> <ul style="list-style-type: none"> • Anti-human T-cell lymphotropic virus (HTLV) antibody (HTLV I/II Ab) • Rapid Plasma Reagin (RPR) – syphilis • Cytomegalovirus (CMV) IgG and IgM • West Nile Virus nucleic acid test • T Cruzi antibody • Strongyloides antibody to assess for previous infection or exposure • T-spot to assess for exposure or history of tuberculosis
---	--

¹ Primary team should follow up on all testing and order follow up testing and consults as indicated prior to proceeding to IEC therapy

² Patients with recent travel out of the country should be considered for some/all of these additional tests

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX C: Infection Prophylaxis Considerations for IEC Therapy

Most patients who receive immune effector cells receive a fludarabine-based chemotherapy regimen prior to IEC infusion. Fludarabine causes immunosuppression and can increase the risk of opportunistic infections. Additionally, patients who receive IEC therapies targeting B-cells are at increased risk of infection due to B-cell aplasia.

Prophylaxis	Preferred Medication	Alternative Medication(s)	Start	Stop	Comment
Viral <ul style="list-style-type: none"> • Herpes simplex • Varicella zoster 	Valacyclovir ¹ 15 mg/kg/day (maximum 500 mg) PO daily or divided twice daily	Acyclovir ¹ 30-45 mg/kg/dose (maximum 800 mg/dose) PO twice daily or 5 mg/kg IV every 8 hours	IEC infusion day	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	-
Hepatitis B (only for patients who are positive for HBsAg or HBcAb)	Entecavir ¹ : <ul style="list-style-type: none"> • 2-15 years old: Consult with Pharmacist for dosing • ≥ 16 years old: Entecavir 0.5 mg PO daily 	Tenofovir disoproxil fumarate ¹ ≥ 2 years old and ≥ 10 kg and adolescents: 8 mg/kg/dose (maximum 300 mg/day) PO daily	2 weeks before IEC	12-24 months post IEC	Consider Infectious Disease and/or Hepatology consult if not already following. Monitor HBV DNA PCR once a month while on prophylaxis and for a year after stopping. Consult Infectious Diseases if entecavir cannot be used or if DNA PCR detectable.
Bacterial (if neutropenia with ANC < 1 K/microliter is expected to last ≥ 7 days)	Levofloxacin ¹ <ul style="list-style-type: none"> • < 5 years old: 10 mg/kg/dose (maximum 250 mg/dose) PO/IV every 12 hours • ≥ 5 years old: 10 mg/kg/dose (maximum 500 mg/dose) PO/IV every 24 hours 	Cefpodoxime ^{1,2} 5 mg/kg/dose (maximum 200 mg per dose) PO twice daily Or Ciprofloxacin ¹ <ul style="list-style-type: none"> • 10 mg/kg/dose (maximum 500 mg per dose) PO twice daily • 10 mg/kg/dose (maximum 400 mg per dose) IV twice daily 	IEC infusion day or when ANC ≤ 0.5 K/microliter	Continue until ANC > 0.5 K/microliter for 3 consecutive days without growth factor support (whichever is longer)	Consult Infectious Diseases if patient is allergic to quinolones and cephalosporins

ANC = absolute neutrophil count

¹ Adjust for renal function

² Cefpodoxime does not cover pseudomonas

Continued on next page

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX C: Infection Prophylaxis Considerations for IEC Therapy - *continued*

Most patients who receive immune effector cells receive a fludarabine-based chemotherapy regimen prior to IEC infusion. Fludarabine causes immunosuppression and can increase the risk of opportunistic infections. Additionally, patients who receive IEC therapies targeting B-cells are at increased risk of infection due to B-cell aplasia.

Prophylaxis	Preferred Medication(s)	Alternative Medication(s)	Start	Stop	Comment
Pneumocystis jirovecii	Pentamidine inhaled or IV ¹ within 1 week prior to IEC infusion and then transition to Sulfamethoxazole/trimethoprim (SMZ/TMP; preferred post-IEC infusion) by 3-4 weeks if counts have recovered: 2.5 – 5 mg/kg/dose trimethoprim (maximum 160 mg) PO twice daily given three days per week on consecutive days (i.e. Fri, Sat, Sun) or on alternating days (i.e. Mon, Wed, Fri) Optional dosing for adolescents: SMZ/TMP 1 double strength tab (800/160 mg) PO daily given three days per week on consecutive days (i.e. Fri, Sat, Sun) or on alternating days (i.e. Mon, Wed, Fri)	-	Within 1 week prior to IEC infusion	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	SMZ/TMP also has activity against toxoplasma and nocardia
		Pentamidine inhaled ≥ 5 years old: 300 mg flat dose every 28 days	Within 1 week prior to IEC infusion	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	Albuterol nebulizer premedication encouraged
		Pentamidine ¹ IV 4 mg/kg (maximum 300 mg per dose) every 21 days	Within 1 week prior to IEC infusion	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	Can cause pancreatitis
		Dapsone 2 mg/kg/dose (maximum 100 mg/dose) PO daily	3-4 weeks post IEC infusion	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	Check G6PD level Use caution if patient has sulfa allergy Can cause hemolytic anemia
		Atovaquone • < 4 months old: 30 mg/kg/day (maximum of 1500 mg/day) PO daily • 4-24 months old: 45 mg/kg/day (maximum of 1500 mg/day) PO daily • > 24 months-13 years old: 30 mg/kg/day (maximum 1500 mg/day) PO daily • > 13 years old: 1500 mg PO daily	Within 1 week prior to IEC infusion	At least 1 year post IEC infusion; may stop after 1 year if CD4 count > 200 cells/microliter	Must take with a fatty meal. Also has activity against toxoplasma, but inferior to SMZ/TMP.

¹ Adjust for renal function

Continued on next page

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX C: Infection Prophylaxis Considerations for IEC Therapy - *continued*

Prophylaxis	Preferred Medication	Alternative Medication	Start	Stop	Comment
Fungal (low risk)	Fluconazole ¹ 12 mg/kg/dose (maximum 400 mg/dose) PO or IV daily	Caspofungin 50 mg/m ² /dose IV daily (maximum 50 mg/day) ²	IEC infusion day	Continue until ANC > 0.5 K/microliter for 3 consecutive days without growth factor support	-
Fungal (high risk) ³	Posaconazole ³ <ul style="list-style-type: none"> • ≤ 12 years old: 4 mg/kg/dose of oral suspension PO three times a day • ≥ 13 years old: 200 mg PO (oral suspension) three times a day or 300 mg PO (delayed release tablets) or IV daily <i>*Tablets preferred in those who can swallow tablets*</i> Voriconazole ³ <ul style="list-style-type: none"> • 2-11 years old or < 50 kg: 9 mg/kg/dose PO every 12 hours (maximum 350 mg/dose) or 8 mg/kg/dose IV every 12 hours • ≥ 12 years old and ≥ 50 kg: 200 mg PO every 12 hours or 4mg/kg/dose IV every 12 hours <i>*Pharmacokinetic monitoring of levels is recommended for both posaconazole and voriconazole*</i>	Caspofungin 50 mg/m ² /dose IV daily (maximum 50 mg/day) ²	IEC infusion day or when high-risk criteria are met	Continue as clinically indicated ³	Posaconazole suspension must be taken with a fatty meal Posaconazole and voriconazole drug-drug interactions via CYP3A4 exist

¹ Adjust for renal function

² Loading dose of antifungals is not needed if it is being used for prophylaxis

³ Posaconazole prophylaxis is recommended for HIGH RISK patients with leukemia, recent allogenic stem cell transplant, prior history of mold infection, neutropenia lasting ≥ 14 days, Grade 3 or 4 CRS/ICANS, those who receive ≥ 3 days of corticosteroids, or those who develop hemophagocytic lymphohistiocytosis (HLH) (see [Appendix N](#)). If corticosteroids are given, continue posaconazole for at least 1 month AFTER COMPLETION of corticosteroids. Do not stop posaconazole or voriconazole prophylaxis if ANC < 1 K/microliter. Isavuconazole may be used if the patient had previously been taking it or if posaconazole is not covered by insurance. In the event posaconazole, voriconazole, or an echinocandin are contraindicated or pose affordability/access issues, then use fluconazole for prophylaxis and consider aspergillus antigen testing at least once a week DURING corticosteroids and for at least a month AFTER completion of corticosteroids. Patients not meeting high risk definitions will be considered to be at LOW RISK for fungal infections and receive prophylaxis as detailed above.

Continued on next page

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX C: Infection Prophylaxis Considerations for IEC Therapy - *continued*

Prophylaxis	Comment
Human Immunodeficiency Virus (HIV)	Antiretroviral Therapy (ART) and monitoring per ID recommendations. Obtain an ID consult on any patient with HIV.
Herpesvirus 6 (HHV-6)	Monitor HHV-6 by quantitative PCR from blood plasma once weekly if neutropenia lasts \geq 14 days, if patient experiences Grade 3 or 4 CRS/ICANS, if patient receives \geq 3 days of corticosteroids, or if patient develops HLH ¹ . HHV-6 monitoring is recommended for at least 30 days after completion of corticosteroids.
CMV	Routine CMV prophylaxis is not required but CMV monitoring by PCR is recommended 1-2 times a week if neutropenia lasts \geq 14 days, if patient experiences Grade 3 or 4 CRS/ICANS, if patient receives \geq 3 days of corticosteroids, or if patient develops hemophagocytic lymphohistiocytosis (HLH) ¹ . CMV monitoring is recommended for at least 30 days after completion of corticosteroids.
Immunoglobulin replacement therapy	Hypogammaglobulinemia may be observed after IEC therapies that target B-cells and IgG levels should be checked in such patients when they develop respiratory infections. Immunoglobulin replacement therapy and/or prophylaxis is only indicated for patients who develop hypogammaglobulinemia (< 500 mg/dL)
Prolonged cytopenias	Grade 3 or 4 cytopenias lasting beyond day 30 have been reported in approximately 30% of patients after IEC therapies. Cytopenias may be managed with filgrastim products; monitor blood counts at least weekly. Continue appropriate prophylactic antimicrobials as described above. Diagnostic bone marrow may be performed to rule out other causes such as myelodysplasia, malignancy, HLH ¹ , or infection.

¹ See [Appendix N](#) for Diagnostic Criteria for IEC-associated Fulminant Hemophagocytic Lymphohistiocytosis (HLH) or Macrophage Activation Syndrome (MAS)

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX D: ASTCT Grading for CRS¹ (Note: CRS grade should be determined at least twice daily and any time there is a change in patient’s status)

CRS Parameter	CRS Grade 1	CRS Grade 2	CRS Grade 3	CRS Grade 4
Fever²	Yes	Yes	Yes	Yes
		With		
Hypotension³	No	Requiring IV fluids but not requiring vasopressors	Requiring one vasopressor with or without vasopressin	Requiring multiple vasopressors (excluding vasopressin)
		And/Or		
Hypoxia³	No	Requiring low-flow O ₂ via nasal cannula ⁴ or blow-by	Requiring O ₂ via high-flow nasal cannula ⁴ , facemask, non-rebreather mask, or Venturi mask	Requiring O ₂ via positive pressure (e.g., CPAP, BiPAP, and mechanical ventilation)

CPAP = continuous positive airway pressure

BiPAP = bilevel positive airway pressure

¹ Organ toxicities associated with CRS may be graded according to CTCAE v5.0 but they do not influence CRS grading

² Fever is defined as temperature ≥ 38°C not attributable to any other cause. In patients who have CRS then receive antipyretics or anti-cytokine therapy such as tocilizumab or corticosteroids, fever is no longer required to grade subsequent CRS severity. In this case, CRS grading is driven by hypotension and/or hypoxia.

³ CRS grade is determined by the more severe event: hypotension or hypoxia not attributable to any other cause. For example, a patient with temperature of 39.5°C, hypotension requiring one vasopressor and hypoxia requiring low-flow nasal cannula is classified as having Grade 3 CRS.

⁴ Low-flow nasal cannula is defined as oxygen (O₂) delivered at ≤ 5 liters/minute. Low flow also includes blow-by oxygen delivery, sometimes used in pediatrics.

High-flow nasal cannula is defined as oxygen delivered at > 5 liters/minute.

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX E: ASTCT Grading of ICANS¹

Symptom/Sign	Grade 1	Grade 2	Grade 3	Grade 4
Neurological assessment (ICE) Score²	7-9	3-6	0 ³ -2	0 ³ (patient is unarousable and unable to perform ICE)
CAPD⁴ score	1-8	1-8	≥ 9 OR Unable to perform CAPD	Critical (obtunded; convulsive status epilepticus; motor weakness; papilledema; CSF OP ≥ 20 mmHg; cerebral edema)
Depressed level of consciousness⁵	Awakens spontaneously	Awakens to voice	Awakens only to tactile stimulus	Patient is unarousable or requires vigorous or repetitive tactile stimuli to arouse; stupor or coma
Seizure	-	-	Any clinical seizure (focal or generalized) that resolves rapidly (< 5 minutes) or non-convulsive seizures on EEG that resolve with intervention	Life-threatening prolonged seizure (≥ 5 minutes) or repetitive clinical or electrical seizures without return to baseline in between
Motor findings⁶	-	-	-	Deep focal motor weakness such as hemiparesis or paraparesis
Raised intracranial pressure⁷ / cerebral edema	-	-	Focal/local edema on neuroimaging ⁸	Diffuse cerebral edema on neuroimaging, decerebrate or decorticate posturing, cranial nerve VI palsy, papilledema, or Cushing's triad

CAPD = Cornell Assessment of Pediatric Delirium

EEG = electroencephalogram

¹ ICANS grade is determined by the most severe event (CAPD and/or ICE score, level of consciousness, seizure, motor findings, raised intracranial pressure/cerebral edema) not attributable to any other cause). For example, a patient with a ICE score of 3 who has a generalized seizure is classified as having Grade 3 ICANS.

² See [Appendix F](#) for Immune Effector Cell –associated Encephalopathy (ICE) Score for patients ≥ 12 years of age

³ A patient with an ICE score of 0 may be classified as having Grade 3 ICANS if the patient is awake with global aphasia or Grade 4 ICANS if the patient is unarousable

⁴ CAPD is intended for patients on Pediatric Service who are ≤ 12 years of age and/or for children > 12 years of age for whom this is developmentally appropriate (see [Appendix G](#))

⁵ Depressed level of consciousness should not be attributable to any other cause (e.g., sedating medication)

⁶ Tremors and myoclonus associated with immune effector cell therapies may be graded according to CTCAE v5.0 but they do not influence ICANS grading

⁷ Ophthalmology may be consulted to assess for papilledema if concern for elevated ICP, but otherwise not needed for all patients

⁸ Intracranial hemorrhage with or without associated edema is not considered a neurotoxicity feature and is excluded from ICANS grading. It may be graded according to CTCAE v5.0

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: *This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.*

APPENDIX F: Immune Effector Cell-associated Encephalopathy (ICE) Score - for patients ≥ 12 years of age

- **Orientation:** Orientation to year, month, city, hospital: 4 points (1 point each)
- **Naming:** Name 3 objects (*e.g.*, clock, pen, button): 3 points (1 point each)
- **Following commands:** (*e.g.*, Show me 2 fingers or close your eyes and stick out your tongue): 1 point
- **Writing:** Ability to write a standard sentence (*e.g.*, Our national bird is the bald eagle): 1 point
- **Attention:** Count backwards from 100 by 10: 1 point

Score 10: No impairment

Score 7-9: Grade 1 ICANS

Score 3-6: Grade 2 ICANS

Score 0-2: Grade 3¹ ICANS

Score 0 due to patient unarousable and unable to perform ICE assessment: Grade 4 ICANS

¹ A patient with an ICE score of 0 may be classified as having Grade 3 ICANS if the patient is awake with global aphasia or may be classified as having Grade 4 ICANS if the patient is unarousable

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX G: Cornell Assessment of Pediatric Delirium (CAPD)¹ score (for children ≤ 12 years of age and/or for children > 12 years of age for whom this is developmentally appropriate)

RASS Score ² (if -4 or -5, do not proceed)											
Answer the following questions based on your interactions with the patient over the course of your shift ³											
	Never 4	Rarely 3	Sometimes 2	Often 1	Always 0		Never 0	Rarely 1	Sometimes 2	Often 3	Always 4
1. Does the child make eye contact with the caregiver?						5. Is the child restless?					
2. Are the child’s actions purposeful?						6. Is the child inconsolable?					
3. Is the child aware of his/her surroundings?						7. Is the child underactive – very little movement while awake?					
4. Does the child communicate needs and wants?						8. Does it take the child a long time to respond to interactions?					

¹ An increasing CAPD score (even if less than 9) may indicate concern for delirium and warrants more frequent monitoring.

²Richmond Agitation Sedation Scale (RASS):

- +4 Combative - Overly combative, violent, immediate danger to staff
- +3 Very Agitated – Pulls or removes tube(s) or catheter(s); aggressive
- +2 Agitated – Frequent non-purposeful movement, fights ventilator
- +1 Restless – Anxious, but movements not aggressive or vigorous
- 0 Alert and Calm
- 1 Drowsy – Not fully alert, but has sustained awakening (eye-opening/eye contact) to voice (≥ 10 seconds)
- 2 Light sedation – Briefly awakens with eye contact to voice (< 10 seconds)
- 3 Moderate sedation – Movement
- 4 Deep sedation – No response to voice, but movement or eye opening to physical stimulation
- 5 Unarousable

³For patients age 1-2 years, the following serve as guidelines to the corresponding questions (1-8):

1. Holds gaze. Prefers primary parent. Looks at speaker.
2. Reaches and manipulates objects, tries to change position, if mobile may try to get up
3. Prefers primary parent, upset when separated from preferred caregivers. Comforted by familiar objects (*i.e.*, blanket or stuffed animal)
4. Uses single words or signs
5. No sustained calm state
6. Not soothed by usual comforting actions, for example, singing, holding, talking, and reading
7. Little if any play, efforts to sit up, pull up, and if mobile crawl or walk around
8. Not following simple directions. If verbal, not engaging in simple dialogue with words or jargon

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX H: Management of CRS

CRS Grade	CRS Parameter	Management		
		Diagnostic Work-Up	Supportive Care	Anti-IEC Therapies
Grade 1	Fever	<ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography Cardiac telemetry and pulse oximetry 	<ul style="list-style-type: none"> Acetaminophen and hypothermia blanket as needed for the treatment of fever Ibuprofen if fever is not controlled with above; use with caution or avoid with thrombocytopenia or renal dysfunction Empiric broad-spectrum antibiotics and consider filgrastim products if neutropenic Maintenance IV fluids for hydration Symptomatic management of constitutional symptoms and organ toxicities as per standard guidelines If not on seizure prophylaxis, initiate levetiracetam 10 mg/kg (maximum 500 mg) PO or IV twice daily 	Consider tocilizumab ¹ for 1 dose for persistent fever lasting greater than 3 days
		Grade 2	Hypotension ²	<ul style="list-style-type: none"> Cardiac telemetry Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography
	Hypoxia			<ul style="list-style-type: none"> Pulse oximetry Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography

PICS = Pediatric Intensive Care

¹See Appendix J for Interleukin-6 Antagonist and Alternative Agents

²Hypotension defined as:

- Age 1-10 years: SBP < [70 + (2 x age in years)] mmHg
- Age > 10 years: SBP < 90 mmHg

Continued on next page

Department of Clinical Effectiveness V4

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX H: Management of CRS - *continued*

CRS Grade	CRS Parameter	Management		
		Diagnostic Work-Up	Supportive Care	Anti-IEC Therapies
Grade 3	Hypotension¹	<ul style="list-style-type: none"> Obtain ECHO if not performed already Cardiac telemetry Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography 	<ul style="list-style-type: none"> Transfer patient to PICS IV fluid boluses as needed as in Grade 2 CRS Use vasopressors as needed Symptomatic management of fever as in Grade 1 CRS Symptomatic management of constitutional symptoms and organ toxicities as per standard guidelines 	<ul style="list-style-type: none"> Tocilizumab² as in Grade 2 CRS if not administered previously; tocilizumab may be repeated every 8 hours for up to 3 doses in a 24-hour period Consider stress dose hydrocortisone 12.5-25 mg/m²/day divided every 6 hours if not already given If still hypotensive on vasopressor and hydrocortisone then consider methylprednisolone or dexamethasone as follows: <ul style="list-style-type: none"> If on one vasopressor: methylprednisolone 1-2 mg/kg/day divided every 6 to 12 hours [or dexamethasone 0.5 mg/kg/dose (maximum 10 mg) IV every 6 hours] If on two vasopressors: methylprednisolone 1 – 2 mg/kg/day divided every 6 to 12 hours (or dexamethasone 1 mg/kg/dose [maximum 20 mg] IV every 6 hours) If vasopressin and norepinephrine equivalent³ is ≥ 15 mcg/minute, follow as in Grade 4 CRS Once CRS improves to Grade 1 or less, taper and/or stop corticosteroids depending on clinical situation
	Hypoxia	<ul style="list-style-type: none"> Pulse oximetry Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography 	<ul style="list-style-type: none"> Supplemental oxygen including high- flow nasal cannula, face mask, non-rebreather mask, or Venturi mask as needed Symptomatic management of fever as in Grade 1 CRS Symptomatic management of constitutional symptoms and organ toxicities as per standard guidelines 	<ul style="list-style-type: none"> Tocilizumab² and methylprednisolone 1 – 2 mg/kg/day divided every 6 to 12 hours (or dexamethasone 0.5 mg/kg/dose [maximum 10 mg] IV every 6 hours) if not administered previously; tocilizumab may be repeated every 8 hours for up to 3 doses in a 24-hour period If there is no improvement in hypoxia within 24 hours or there is rapid progression of pulmonary infiltrates or sharp increase in FiO₂ requirements, methylprednisolone 1-2 mg/kg/day divided every 6 to 12 hours [or dexamethasone 1 mg/kg/dose (maximum 20 mg) IV every 6 hours] Once CRS improves to Grade 1 or less, taper and/or stop corticosteroids depending on clinical situation

¹Consider stress-dose hydrocortisone for patients with vasopressor-resistant hypotension attributed to adrenal insufficiency

²See [Appendix J](#) for Dosing of IL-6 Antagonists and Alternative Agents

³VASST Trial vasopressor equivalent equation: norepinephrine equivalent dose = [norepinephrine (mcg/minute)] + [dopamine (mcg/kg/minute) / 2] + [epinephrine (mcg/minute)] + [phenylephrine (mcg/minute) / 10]

Continued on next page

Department of Clinical Effectiveness V4

Approved by the Executive Committee of the Medical Staff on 09/15/2020

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX H: Management of CRS - *continued*

CRS Grade	CRS Parameter	Management		
		Diagnostic Work-Up	Supportive Care	Anti-IEC Therapies
Grade 4	Hypotension	<ul style="list-style-type: none"> Obtain ECHO if not performed already Cardiac telemetry Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography 	<ul style="list-style-type: none"> Transfer patient to PICS IV fluid boluses as needed as in Grade 2 CRS Vasopressors as in Grade 3 CRS Use vasopressors as needed Symptomatic management of fever as in Grade 1 CRS Symptomatic management of constitutional symptoms and organ toxicities as per standard guidelines 	<ul style="list-style-type: none"> Tocilizumab¹ as in Grade 2 CRS if not administered previously; tocilizumab may be repeated every 8 hours for up to 3 doses in a 24-hour period Consider stress dose hydrocortisone 12.5 – 25 mg/m²/day divided every 6 hours if not already given If still hypotensive on vasopressor and hydrocortisone then consider methylprednisolone as follows: <ul style="list-style-type: none"> Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by rapid taper as per clinical situation If hypotension is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable
	Hypoxia	<ul style="list-style-type: none"> Monitor oxygen saturation while on mechanical ventilation Fever work-up if not previously performed <ul style="list-style-type: none"> Assess for infection with blood and urine cultures, and chest radiography 	<ul style="list-style-type: none"> Transfer patient to PICS Positive pressure ventilation including CPAP, BiPAP, mechanical ventilation Symptomatic management of fever as in Grade 1 CRS Symptomatic management of constitutional symptoms and organ toxicities as per standard guidelines 	<ul style="list-style-type: none"> Tocilizumab¹ as in Grade 2 CRS if not administered previously; tocilizumab may be repeated every 8 hours for up to 3 doses in a 24-hour period Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by rapid taper as per clinical situation If hypoxia is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable

¹ See [Appendix J](#) for Dosing of IL-6 Antagonists and Alternative Agents

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX I: Management of ICANS

ICANS Grade	Sign or symptom	Management		
		Diagnostic Work-up	Supportive Care	Anti-IEC Therapies
Grade 1	Encephalopathy and/or depressed level of consciousness	<ul style="list-style-type: none"> • MRI imaging of the brain with and without contrast; CT of brain without contrast may be performed if MRI is not feasible; MRI spine if focal deficits are noted • Neurology consultation • ICE Score and/or CAPD assessment every 6 hours or more frequently if clinically indicated • EEG • Consider diagnostic lumbar puncture if other causes of encephalopathy are suspected (<i>e.g.</i>, infections, autoimmune, leptomenigeal disease) <ul style="list-style-type: none"> ◦ Add a meningitis-encephalitis panel from CSF in patients with neurologic symptoms that persist or worsen after ICANS therapy and/or if symptoms start after corticosteroids 	<ul style="list-style-type: none"> • Vigilant supportive care; aspiration precautions; IV hydration • Withhold oral intake of food/medications/fluids and assess swallowing; convert all oral medications and/or nutrition to IV if swallowing is impaired • Avoid medications that cause central nervous system depression • Low doses of lorazepam after EEG is performed [0.02 mg/kg/dose (maximum 0.5 mg) IV every 8 hours] or haloperidol [0.01 mg/kg/dose (maximum 0.5 mg) IV every 6 hours] may be used with careful monitoring for agitated patients • If no seizures on EEG, continue prophylactic levetiracetam • If EEG shows focal or generalized convulsive or non-convulsive seizure or convulsive status epilepticus, refer to further management as in Grade 3 or 4 ICANS 	<ul style="list-style-type: none"> • If associated with concurrent CRS, add tocilizumab¹ • If no response to tocilizumab or for isolated ICANS, start dexamethasone 0.5 mg/kg/dose (maximum 10 mg) IV for 1 dose (or methylprednisolone equivalent) and reassess in 6 hours or earlier if clinically indicated
Grade 2	Encephalopathy and/or depressed level of consciousness	<ul style="list-style-type: none"> • Neurological work-up as in Grade 1 ICANS 	<ul style="list-style-type: none"> • Supportive care as in Grade 1 ICANS 	<ul style="list-style-type: none"> • Dexamethasone 0.5 mg/kg/dose (maximum 10 mg) IV every 12 hours (or methylprednisolone equivalent) <ul style="list-style-type: none"> ◦ If associated with concurrent CRS, add tocilizumab¹ • Once ICANS improves to Grade 1 or less, taper and/or stop corticosteroids depending on clinical situation

¹ See [Appendix J](#) for Dosing of IL-6 Antagonists and Alternative Agents

Continued on next page

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX I: Management of ICANS - *continued*

ICANS Grade	Sign or symptom	Management		
		Diagnostic Work-up	Supportive Care	Anti-IEC Therapies
Grade 3	Encephalopathy and/or depressed level of consciousness	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS Consider repeat neuro-imaging (CT or MRI) every 2-3 days for persistent > Grade 3 encephalopathy Consider diagnostic lumbar puncture if Grade 3 encephalopathy persists > 2 days or earlier if other causes are suspected (<i>e.g.</i>, infections, autoimmune, leptomenigeal disease) <ul style="list-style-type: none"> ○ Add a meningitis-encephalitis panel from CSF in patients with neurologic symptoms that persist or worsen after ICANS therapy and/or if symptoms start after corticosteroids. 	<ul style="list-style-type: none"> Supportive care as in Grade 1 ICANS Consider PICS transfer If there are new abnormal findings on brain imaging¹ not related to primary malignancy, control hypertension with the goal of MAP within 20-25 mmHg of baseline MAP; correct any uremia (dialysis if needed) and/or coagulopathy (transfuse to keep platelets > 20-50 K/microliter, fibrinogen > 200 mg/dL and INR < 1.5) 	<ul style="list-style-type: none"> Dexamethasone 0.5 mg/kg/dose (maximum 10 mg) IV every 6 hours (or methylprednisolone equivalent) <ul style="list-style-type: none"> ○ If associated with concurrent CRS, add tocilizumab² If Grade 3 encephalopathy is persistent for > 24 hours, increase dexamethasone to 1 mg/kg/dose (maximum 20 mg) IV every 6 hours (or methylprednisolone equivalent) Once ICANS improves to Grade 1 or less, taper and/or stop corticosteroids depending on clinical situation
	Seizure	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS EEG if clinically indicated (<i>e.g.</i>, ongoing seizures, depressed level of consciousness) Rule out other potential causes of seizure (<i>i.e.</i>, beta-lactams, etc.) 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS For focal or generalized convulsive seizures, or non-convulsive seizures, treat as per Appendix K 	<ul style="list-style-type: none"> Dexamethasone 1 mg/kg/dose (maximum 20 mg) IV every 6 hours (or methylprednisolone equivalent) <ul style="list-style-type: none"> ○ If associated with concurrent CRS, add tocilizumab² Once ICANS improves to Grade 1 or less, taper and/or stop corticosteroids depending on clinical situation
	Focal cerebral edema	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS Consider repeat neuro-imaging (CT or MRI) every 24 hours until edema resolves or more frequently if clinically indicated 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS 	<ul style="list-style-type: none"> If focal edema is in brain stem or thalamus, methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by taper depending on clinical situation <ul style="list-style-type: none"> ○ If associated with concurrent CRS, add tocilizumab² If focal edema is in other areas of brain, methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 1 day; assess daily and continue or taper depending on clinical situation <ul style="list-style-type: none"> ○ If associated with concurrent CRS, add tocilizumab²

¹Abnormal findings on imaging where correction of hypertension, uremia, and/or coagulopathy should be performed include changes suggestive of typical or atypical posterior reversible encephalopathy syndrome (PRES), temporal lobe and limbic system encephalitis (autoimmune or infection), acute disseminated encephalomyelitis, emboli, vasculitis, strokes, and/or seizure-related changes

²See [Appendix J](#) for Dosing of IL-6 Antagonists and Alternative Agents

Continued on next page

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX I: Management of ICANS - *continued*

ICANS Grade	Sign or symptom	Management		
		Diagnostic Work-Up	Supportive Care	Anti-IEC Therapies
Grade 4	Encephalopathy and/or depressed level of consciousness	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS Repeat neuro-imaging and lumbar puncture as in Grade 3 ICANS 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS Consider mechanical ventilation for airway protection If there are new abnormal findings on brain imaging¹ not related to primary malignancy, control hypertension with the goal of maintaining MAP within 20-25 mmHg of baseline MAP; correct any uremia (dialysis if needed) and/or coagulopathy (transfuse to keep platelets > 20 - 50 K/microliter, fibrinogen > 200 mg/dL and INR < 1.5) 	<ul style="list-style-type: none"> Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by taper as clinically indicated; if associated with concurrent CRS, add tocilizumab² Continue corticosteroids until improvement to less than or equal to Grade 1 ICANS and then taper and stop corticosteroids depending on clinical situation If Grade 4 ICANS is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable
	Seizure	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS Rule out other potential causes of seizure (i.e., beta-lactams, etc.) 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS For focal or generalized convulsive or non-convulsive seizure or convulsive status epilepticus, treat as in Appendix K For convulsive status epilepticus, treat as in Appendix L 	<ul style="list-style-type: none"> Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by taper as clinically indicated; if associated with concurrent CRS, add tocilizumab² If Grade 4 ICANS is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable
	Motor Weakness	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS MRI with and without contrast of the spine 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS 	<ul style="list-style-type: none"> Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by taper as clinically indicated; if associated with concurrent CRS, add tocilizumab² If Grade 4 ICANS is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable
	Diffuse cerebral edema or raised intracranial pressure	<ul style="list-style-type: none"> Neurological work-up as in Grade 1 ICANS Consider repeat neuro-imaging as in focal cerebral edema from Grade 3 ICANS 	<ul style="list-style-type: none"> Transfer to PICS Supportive care as in Grade 1 ICANS For diffuse cerebral edema or signs of raised intracranial pressure, treat as in Appendix M 	<ul style="list-style-type: none"> Methylprednisolone 30 mg/kg/day (maximum 1,000 mg/day) in divided doses IV for 3 days followed by taper as clinically indicated; if associated with concurrent CRS, add tocilizumab² If Grade 4 ICANS is refractory for > 24 hours or if patient is deteriorating rapidly, consider additional therapies (see Appendix J) including activation of safety switches if applicable

¹Abnormal findings on imaging where correction of hypertension, uremia, and/or coagulopathy should be performed include changes suggestive of typical or atypical posterior reversible encephalopathy syndrome (PRES), temporal lobe and limbic system encephalitis (autoimmune or infection), acute disseminated encephalomyelitis, emboli, vasculitis, strokes, and/or seizure-related changes

²See [Appendix J](#) for Dosing of IL-6 Antagonists and Alternative Agents

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX J: Recommendations for Use of IL-6 Antagonists and Alternative Agents for Management of CRS and ICANS

Drug	Recommended Dose for CRS and/or ICANS	Maximum Dose	Mechanism of Action	Comments
Tocilizumab ¹	Less than 30 kg: 12 mg/kg IV for up to three doses in a 24-hour period 30 kg and greater: 8 mg/kg IV for up to three doses in a 24-hour period	Maximum 800 mg per dose	IL-6 receptor antagonist	<ul style="list-style-type: none"> • Maximum of 4 doses total over the entire course of CRS and ICANS • Dose may be repeated every 8 hours for up to three doses in a 24-hour period
Siltuximab ²	11 mg/kg IV once	No more than 1 dose in a 3-week period	Binds to both soluble and membrane bound IL-6 Neutralizes IL-6	<ul style="list-style-type: none"> • Recommended primarily for patients who are intolerant to tocilizumab • No more than 1 dose in a 3 week period
Anakinra ³	Children ≥ 2 years and Adolescents: 1 mg/kg (maximum 100 mg) subcutaneously daily for 7 days	-	IL-1 receptor antagonist	<ul style="list-style-type: none"> • Renal dose adjustment may be needed for creatinine clearance < 30 mL/minute
Cyclophosphamide	1,500 mg/m ² IV for one dose	-	Alkylating agent	<ul style="list-style-type: none"> • Give with mesna 1500 mg/m² IV over 24 hours for one dose
Anti-thymocyte globulin (rabbit)	1-2 mg/kg IV daily for 3 days	-	Immunosuppressant	<ul style="list-style-type: none"> • Hypersensitivity reactions can occur; premedicate with diphenhydramine and scheduled dose of corticosteroid • Infuse over a minimum of 6 hours
Safety switches	-	-	-	<ul style="list-style-type: none"> • If the IEC product contains a safety switch (e.g., iCaspase-9 or EGFRt-positive), the corresponding drug to eliminate those cells can be considered in doses according to manufacturer • Examples include rimiducid to eliminate iCaspase-9 or cetuximab to eliminate EGFRt-positive cells

¹ MD Anderson formulary restricted for use in CRS/ICANS and for use in hemophagocytic lymphohistiocytosis (HLH), see [Appendix N](#)

² MD Anderson formulary restricted for use in CRS/ICANS

³ Not on MD Anderson formulary; use at MD Anderson is based on internal data in patients with tocilizumab and/or siltuximab failure

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX K: Management of Focal or Generalized Convulsive or Non-Convulsive Seizures

- Assess CAB / consider airway protection / check blood glucose
- Consult Neurology
- For focal and generalized convulsive seizures, lorazepam 0.05 mg/kg (maximum 1 mg) IV; repeat dose every 5 minutes (to a maximum 4 doses) to control electrographical seizures
- Levetiracetam 40 mg/kg (maximum 2,500 mg) IV bolus (in addition to maintenance dose)
- Replete with magnesium as needed to maintain magnesium level > 2 mg/dL
- Thiamine 100 mg IV every 8 hours for 5 days
- If non-convulsive seizures persist, transfer to PICS and add phenobarbital loading dose of 10-20 mg/kg (maximum 1,000 mg) IV (monitor for respiratory depression, bradycardia and hypotension)
- Maintenance doses after resolution of non-convulsive status epilepticus
 - Lorazepam 0.05 mg/kg (maximum 1 mg) IV every 8 hours for 3 doses
 - Levetiracetam 15 mg/kg (maximum 1,500 mg) IV every 12 hours
 - Phenobarbital 1-3 mg/kg IV every 12 hours
 - Monitor for respiratory depression, bradycardia and hypotension
 - Assess for drug-drug interactions (*i.e.*, may induce metabolism of azole antifungals or other CYP3A4 substrates) and consider alternative therapy if drug interactions are significant
 - Target serum trough levels 15-40 mcg/mL

APPENDIX L: Management of Convulsive Status Epilepticus

- Assess CAB / airway protection / high flow O₂, check blood glucose
- Transfer to PICS
- Consult Neurology
- Lorazepam 0.1 mg/kg (maximum 4 mg) IV; repeat dose after at least 1 minute (to a maximum of 2 doses) to control seizures
- Levetiracetam 40 mg/kg (maximum 2,500 mg) IV bolus (in addition to maintenance dose)
- Replete with magnesium as needed to maintain magnesium level > 2 mg/dL
- Thiamine 100 mg IV every 8 hours for 5 days
- If seizures persist, add phenobarbital at a loading dose 10-20 mg/kg (maximum 1,000 mg) IV (monitor for respiratory depression, bradycardia and hypotension)
- If refractory, consider additional therapies (see [Appendix J](#)) including activation of safety switches if applicable
- Maintenance doses after resolution of status epilepticus
 - Lorazepam 0.05 mg/kg (maximum 1 mg) IV every 8 hours for 3 doses
 - Levetiracetam 30 mg/kg (maximum 1,500 mg) IV every 12 hours or increase the current dose by 10 mg/kg IV every 12 hours
 - Phenobarbital 1-3 mg/kg IV every 12 hours
 - Monitor for respiratory depression, bradycardia and hypotension
 - Assess for drug-drug interactions (*i.e.*, may induce metabolism of azole antifungals or other CYP3A4 substrates) and consider alternative therapy if drug interactions are significant
 - Target serum trough levels 15-40 mcg/mL
 - Continuous EEG if seizures are refractory to treatment

CAB = circulation, airway, breathing

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX M: Management of Diffuse Cerebral Edema and/or Raised Intracranial Pressure

<p>For papilledema without diffuse cerebral edema or other signs of raised intracranial pressure</p>	<ul style="list-style-type: none"> • Acetazolamide 15 mg/kg (maximum 1,000 mg) IV followed by 8-12 mg/kg (maximum 1,000 mg) IV every 12 hours (monitor renal function and acid/base balance once or twice daily and adjust dose accordingly) • Dexamethasone 1 mg/kg/dose (maximum 20 mg) IV every 6 hours (or methylprednisolone equivalent) and start taper after resolution of papilledema
<p>For diffuse cerebral edema on neuroimaging or signs of raised intracranial pressure such as decerebrate or decorticate posturing, cranial nerve VI palsy, or Cushing’s triad</p>	<ul style="list-style-type: none"> • Methylprednisolone 30 mg/kg/day in divided doses IV for 3 days followed by taper as clinically indicated • Elevate head end of patient’s bed to an angle of 30 degrees • Hyperventilation to achieve target PaCO₂ of 28-30 mmHg, but maintained for no longer than 24 hours • Hyperosmolar therapy with either mannitol 20% or hypertonic saline (3% or 23.4% as detailed below) <ul style="list-style-type: none"> ◦ Mannitol: initial dose 0.5-1 g/kg IV; maintenance dose 0.25-1 g/kg IV every 6 hours while monitoring metabolic profile and serum osmolality every 6 hours; and withhold mannitol infusion if serum osmolality is ≥ 320 mOsm/kg or osmolality gap is ≥ 40) ◦ Hypertonic 3% saline: initial dose 5 mL/kg IV over 15 minutes, maintenance dose of 1 mL/kg/hour IV to reach a target sodium level of 150-155 mEq/L (monitor electrolytes every 4 hours; hold infusion if serum sodium levels reach > 155 mEq/L) ◦ Hypertonic 23.4% saline (for patients with imminent herniation): 0.5 – 1 mL/kg. Dose to be administered by physician; repeat after 15 minutes, if needed • If patient has ommaya reservoir, drain CSF to target OP < 20 mmHg • Control hypertension with the goal of maintaining mean arterial pressure (MAP) within 20-25 mmHg of baseline MAP; correct any uremia (dialysis if needed) and/or coagulopathy (transfuse to keep platelets > 20-50 K/microliter, fibrinogen > 200 mg/dL and INR < 1.5) • Consider neurosurgery consultation and IV anesthetics for burst-suppression pattern on EEG; transfuse to keep platelets ≥ 100 K/microliter if possible and correct coagulopathy in case of surgical intervention • Consider additional therapies (see Appendix J) including activation of safety switches if applicable • Metabolic profile every 6 hours and daily CT scans of head without contrast, with adjustments in usage of aforementioned medications to prevent rebound cerebral edema, renal failure, electrolyte abnormalities, hypovolemia and hypotension

OP = opening pressure


Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX N: Diagnostic Criteria for IEC-Related Hemophagocytic Lymphohistiocytosis (HLH) or Macrophage Activation Syndrome (MAS)

- Consider HLH/MAS if a patient has a peak ferritin > 10,000 ng/mL during the CRS phase and develops any two of the following organ toxicities after IEC therapy
 - ≥ Grade 3 increase in bilirubin, aspartate transaminase, or alanine transaminase¹
 - ≥ Grade 3 oliguria or increase in creatinine¹
 - ≥ Grade 3 pulmonary edema¹
 - Presence of hemophagocytosis by morphology and/or CD68 immunohistochemistry in bone marrow or organs
- If HLH/MAS is suspected, obtain baseline fasting triglyceride level and serum soluble IL-2 receptor

¹Grading as per Common Terminology Criteria for Adverse Events, version 5

MANAGEMENT OF IEC-ASSOCIATED HLH / MAS



²Intrathecal cytarabine

- 1 – 1.99 years: 30 mg
- 2 – 2.99 years: 50 mg
- At least 3 years old: 70 mg

³Intrathecal hydrocortisone

- 1 – 1.99 years: 15 mg
- 2 – 2.99 years: 25 mg
- At least 3 years old: 50 mg

IEC Therapy Toxicity Assessment and Management (also known as CARTOX) – Pediatric

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX O: Determine if the Subject Has Allogeneic IEC-associated Acute Graft-Versus-Host Disease (GVHD)

If a subject has any of the following symptoms or signs within the first 3 months after allogeneic IEC therapy, the subject may have acute GVHD if the symptoms or signs are not attributable to any other cause.

1. Skin rash
2. Diarrhea (may also be associated with nausea, vomiting, and/or anorexia due to upper GI GVHD)¹
3. Total bilirubin \geq 2 mg/dL

¹ Isolated upper GI GVHD with nausea, vomiting, and/or anorexia is uncommon following allogeneic lymphocyte infusion

APPENDIX P: Determine the Grade of IEC-associated Acute GVHD

GVHD Target Organ Staging

Stage	Skin	Liver (bilirubin)	Lower GI (stool output/day)
0	No active (erythematous) GVHD rash	< 2 mg/dL	< 10 mL/kg/day or < 4 episodes/day
1	Maculopapular rash < 25% BSA	2 - 3 mg/dL	10-19.9 mL/kg/day or 4-6 episodes/day
2	Maculopapular rash 25 – 50% BSA	3.1 - 6 mg/dL	20-30 mL/kg/day or 7-10 episodes/day
3	Maculopapular rash > 50% BSA	6.1 - 15 mg/dL	> 30 mL/kg/day or > 10 episodes/day
4	Generalized erythroderma (> 50% BSA) plus bullous formation and desquamation > 5% BSA	> 15 mg/dL	Severe abdominal pain with or without ileus or grossly bloody stool (regardless of stool volume)

BSA = body surface area

Overall Clinical Grade (based on most severe organ involvement)

Grade	Comment
0	No stage 1-4 of any organ
I	Stage 1-2 skin without liver, upper GI ² , or lower GI involvement
II	Stage 3 rash and/or stage 1 liver and/or stage 1 upper GI ² and/or stage 1 lower GI
III	Stage 2-3 liver and/or stage 2-3 lower GI, with stage 0-3 skin and/or stage 0-1 upper GI ²
IV	Stage 4 skin, liver, or lower GI involvement, with stage 0-1 upper GI ²

² Isolated upper GI GVHD with nausea, vomiting, and/or anorexia is uncommon following allogeneic lymphocyte infusion

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson’s specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient’s care. This algorithm should not be used to treat pregnant women.

APPENDIX Q: Manage IEC-associated Acute GVHD

Grade	Sign or Symptom	Management
Grade I	Skin rash	<ul style="list-style-type: none"> • Skin biopsy, preferably non-sun exposed site • Hydrocortisone cream 1% twice daily to face • Triamcinolone cream 0.1% three times daily to affected body area • If patient fails triamcinolone, may consider clobetasol cream 0.05% twice daily to body; limit use to no longer than 1-2 weeks • All corticosteroid creams should be followed by an emollient such as CeraVe, Aquaphor or Eucerin (creams not lotions) 20-40 minutes after application of corticosteroid
Grade II-IV	<ul style="list-style-type: none"> • Skin rash > 50% BSA and/or • Total bilirubin > 2 mg/dL and/or • Diarrhea > 10 mL/kg/day 	<ul style="list-style-type: none"> • Skin biopsy as above for rash • Gastrointestinal consult for flexible sigmoidoscopy with or without upper GI endoscopy with duodenal biopsy¹ • DO NOT give GI prep (GoLytely, etc.) unless full colonoscopy ordered • Stool culture for <i>C. difficile</i> and GI multiplex panel • DO NOT wait for completion of these procedures to start systemic therapy • Start prednisone 2 mg/kg/day orally or methylprednisolone equivalent in divided doses

BSA = body surface area

¹Upper GI endoscopy may be considered if patient has nausea, vomiting, and/or anorexia and upper GI GVHD is suspected

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

SUGGESTED READINGS

- Fattal-Valevski, A., Bloch-Mimouni, A., Kivity, S., Heyman, E., Brezner, A., Strausberg, R., ... Goldberg-Stern, H. (2009). Epilepsy in children with infantile thiamine deficiency. *Neurology*, 73(11), 828-33. <https://doi.org/10.1212/WNL.0b013e3181b121f5>.
- Frisén, L. (1982). Swelling of the optic nerve head: a staging scheme. *Journal of Neurology, Neurosurgery & Psychiatry*, 45(1), 13-18. <https://doi.org/10.1136/jnnp.45.1.13>
- Giavridis, T., van der Stegen, S., Eyquem, J., Hamieh, M., Piersigili, A., & Sadelain, M. (2018). CAR T cell-induced cytokine release syndrome is mediated by macrophages and abated by IL-1 blockade. *Nature Medicine*, 24, 731-738. <https://doi.org/10.1038/s41591-018-0041-7>
- Kelly, A. & Ramanan, A. (2008). A case of macrophage activation syndrome successfully treated with anakinra. *Nature Clinical Practice Rheumatology*, 4(11), 615-20. <http://www.ncbi.nlm.nih.gov/pubmed/18825135>
- Henter, J., Horne, A., Aricó, M., Egeler, R. M., Filipovich, A. H., Imashuku, S., ... Janka, G for the Histocyte Society. (2007). HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatric Blood and Cancer*, 48(2), 124-131. <https://doi.org/10.1002/pbc.21039>
- Lee, D. W., Gardner, R., Porter, D. L., Louis, C. U., Ahmed, N., Jensen, M., ... Mackall, C. L. (2014). Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*, 124(2), 188-195. <https://doi.org/10.1182/blood-2014-05-552729>
- Lee, D., Santomasso, B., Locke, F., Ghobadi, A., Turtle, C., Brudno, J., ... Neelapu, S. (2019). ASTCT consensus grading for cytokine release syndrome and neurological toxicity associated with immune effector cells. *Biology of Blood and Marrow Transplantation*, 25(4), 625-638. <https://doi.org/10.1016/j.bbmt.2018.12.758>
- Mahadeo, K., Khazal, S., Abdel-Azim, H., Fitzgerald, J., Taraseviciute, A., Bollard, B., ... Pediatric Acute Lung Injury and Sepsis Investigators (PALISI) Network. (2019). Management guidelines for paediatric patients receiving chimeric antigen receptor T cell therapy. *Nat Rev Clin Oncol*, 16(1), 45-63. <https://doi.org/10.1038/s41571-018-0075-2>.
- Neelapu, S. S., Tummala, S., Kebriaei, P., Wierda, W., Gutierrez, C., Locke, F. L., ... Westin, J. (2018). Chimeric antigen receptor T-cell therapy—assessment and management of toxicities. *Nature Reviews Clinical Oncology*, 15(1), 47-62. <https://doi.org/10.1038/nrclinonc.2017.148>
- Norelli, M., Camisa, B., Barbiera, G., Falcone, L., Purevdorj, A., Genua, M., ... Bondanza, A. (2018). Monocyte-derived IL-1 and IL-6 are differentially required for cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nature Medicine*, 24, 739-748. <https://doi.org/10.1038/s41591-018-0036-4>
- Pearl, P. (2016). Amenable Treatable Severe Pediatric Epilepsies. *Semin Pediatr Neurol*, 23(2), 158-66. <https://doi.org/10.1016/j.spen.2016.06.004>
- Rhodes, A., Evans, L., Alhazzani, W., Levy, M., Antonelli, M., Ferrer, R., ... Dellinger, R. (2017). Surviving sepsis campaign: International guidelines for management of sepsis and septic shock: 2016. *Intensive Care Medicine*, 43, 304-377. <https://doi.org/10.1007/s00134-017-4683-6>
- Russell, J. A., Walley, K. R., Singer, J., Gordon, A. C., Hébert, P. C., Cooper, D. J., ... Dieter, A. for the VASST Investigators. (2008). Vasopressin versus norepinephrine infusion in patients with septic shock. *The New England Journal of Medicine*, 358(9), 877-887. <https://doi.org/10.1056/NEJMoa067373>
- Sorrer, M., Logan, B., Zhu, X., Rizzo, J., Cooke, K., McCarthy, P., ... Pasquini, M. (2015). Prospective validation of the predictive power of the hematopoietic cell transplantation comorbidity index: A center for international blood and marrow transplant research study. *Biology of Blood and Marrow Transplantation*, 21(8), 1479-1487. <https://doi.org/10.1016/j.bbmt.2015.04.004>
- The Pediatric Acute Lung Injury Consensus Conference Group. (2015). Pediatric acute respiratory distress syndrome: consensus recommendations from the pediatric acute lung injury consensus conference. *Pediatric Critical Care Medicine*, 16(5), 428-439. <https://doi.org/10.1097/PCC.0000000000000350>
- US Department of Health and Human Services. (2017). Common terminology criteria for adverse events (CTCAE version 5.0). https://ctep.cancer.gov/protocoldevelopment/electronic_applications/docs/CTCAE_v5_Quick_Reference_5x7.pdf

Disclaimer: *This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.*

DEVELOPMENT CREDITS

This practice consensus statement is based on majority opinion of the Pediatric IEC Therapy Toxicity Assessment and Management experts at the University of Texas MD Anderson Cancer Center for the patient population. These experts included:

Linette J. Ewing, MD (Pediatrics – Patient Care)
Alison Gulbis, PharmD (Pharmacy Clinical Programs)[†]
Sajad Khazal, MD (Pediatrics – Patient Care)[†]
Kris M. Mahadeo, MD (Pediatrics – Patient Care)[†]
Rodrigo Mejia, MD (Pediatrics – Patient Care)[†]
Maria Estela Mireles, PharmD (Pharmacy Clinical Programs)[†]
Christina Perez[♦]
Demetrios Petropoulos, MD (Pediatrics – Patient Care)[†]
Shehla Razvi, MD (Pediatrics – Patient Care)[†]
Basirat Shoberu, PharmD (Pediatrics – Patient Care)[†]
Elizabeth Shpall, MD (Stem Cell Transplantation)
John Slopis, MD (Neuro-Oncology)
Milena Zhang, PharmD[♦]

[†] Core Development Team Lead

[♦] Clinical Effectiveness Development Team